Tracheostomy on the intensive care unit for adult patients

B.G. Fikkers¹, in collaboration with the committee "Guidelines Tracheostomy" of the $NVIC^2$

¹ Dr. B.G. Fikkers, Radboud University Nijmegen Medical Centre <u>b.fikkers@ic.umcn.nl</u>

² Drs. P. Breedveld, azM, Maastricht <u>pbre@shee.azm.nl</u> Drs. D. Dongelmans, AMC, Amsterdam <u>d.a.dongelmans@amc.nl</u> Prof. Dr. J.G. van der Hoeven, Radboud University Nijmegen Medical Centre <u>j.vanderhoeven@ic.umcn.nl</u> Dr. M.J. Schultz, St Michaels Hospital University of Toronto, Canada <u>m.j.schultz@amc.nl</u> Dr. J.J. Spijkstra, VU University Medical Center, Amsterdam <u>jj.spijkstra@vumc.nl</u> Drs. R.J. de Wit, Medisch Spectrum Twente, Enschede <u>r.dewit@ziekenhuis-mst.nl</u>

Conflict of interest: None

Financial support: None

Samenvatting

Op de intensive care heeft de percutane tracheostomie de chirurgische tracheostomie verdrongen t.b.v. volwassen patienten. Het is lastig precieze aanbevelingen te geven aangaande de optimale timing van een (percutane) tracheostomie, aangezien dit afhangt van de klinische toestand en de prognose van de patiënt. Bovendien behoren vrijwel alle aanbevelingen tot "level D of E evidence".

Indien de beademing langer lijkt te gaan duren dan twee weken, dan is een tracheostomie geïndiceerd (Level C).

De meeste contra-indicaties voor een percutane tracheostomie zijn relatief en hangen af van lokale richtlijnen en individuele ervaring (Level E).

Geen van de technieken van percutane tracheostomie heeft significante voordelen boven een andere en de keus hangt af van individuele ervaring en voorkeur (Level C).

Summary

Percutaneous tracheostomy has replaced surgical tracheostomy as the preferred technique for adult patients in the intensive care unit (ICU). It is difficult to give clear guidelines on the optimal timing of (percutaneous) tracheostomy, because this depends on the clinical situation and prognosis of the patient. Moreover, most recommendations are based on level D or E evidence.

When ventilation is expected to last longer than two weeks, early percutaneous tracheostomy, compared to prolonged translaryngeal intubation, is to be preferred (Level C).

Most contra-indications for percutaneous tracheostomy are relative and depend on individual experience (Level E).

None of the techniques of percutaneous tracheostomy has significant advantages or disadvantages over another and choice depends mainly on individual preference (Level C).

1) Outline

This guideline describes the indications for, and the procedure of percutaneous tracheostomy (PT) in adult patients in the intensive care unit (ICU). It is based on two published reviews in Dutch journals ^{1;2} and a recent international review ³. The procedures in an emergency situation are briefly described in paragraph 5 e.

In the literature, the terms tracheotomy and tracheostomy are inconsistently applied and most often used interchangeably ⁴. The committee prefers the term tracheostomy, reserving the term tracheotomy only for the act creating an opening in the trachea.

2) Indications

In general, for patients in the ICU, the majority of indications for surgical and PT are identical (Table 1) $^{5;6}$. The percutaneous technique is preferred $^{7;8}$, unless specific contra-indications exist (see below) (Level C).

3) Advantages of tracheostomy

Tracheostomy offers a number of practical advantages, both to the patient as to the medical and nursing staff, compared to endotracheal intubation (Table 2) $^{9;10}$ (Level D).

4) Contra-indications for tracheostomy

Most contra-indications are relative and depend on individual expertise (Table 3) $^{11;12}$. Patient selection is important, in particular when sufficient expertise is not available (Level E).

5) Special subgroups with relative contra-indications

a. Obesitas

Several case reports and one series involving 13 patients were published describing successful accomplishment of PT in morbidly obese patients ¹³. A recent study has presented experiences with PT in 73 morbidly obese patients (from a group of 474 patients). An almost five-fold increase of major complications in the obese patients group compared to the control group was seen ¹⁴. Therefore, in obese patients, the risks and benefits of (percutaneous) tracheostomy should be carefully balanced, and an experienced team should perform the procedure. The routine use of an extended-length tracheostomy tubes should be considered ¹⁵ (Level D).

b. Poststernotomy

Surgical tracheostomies are frequently colonized and infected and therefore constitute a risk factor for mediastinitis after cardiac surgery ¹⁶. In 1973, cricothyrotomy was advocated to prevent median sternotomy infections ¹⁷. More recently it was shown that PT is safe in this respect and the procedure can be advised in early poststernotomy patients who are expected to be ventilated for a prolonged period of time ¹⁸ (Level E).

c. Trauma with suspected neck injury

PT can be safely performed without cervical spine clearance and neck extension in trauma patients who require long-term airway management ^{19;20}. However, because the procedure is more difficult, physicians with limited PT experience should not perform it (Level D).

d. Coagulation abnormalities

Advantages of PT are the tight fit of the tract around the tracheostomy tube, which compresses any small bleeding vessels. Since it involves minimum tissue dissection, it is therefore suitable for patients with a high bleeding risk. Although coagulation abnormalities are no longer an absolute contraindication, correction of haemostasis should be carefully performed ²¹, aiming for levels comparable to those suggested in neuraxial blockade, i.e. INR<1.8, APTT<1.5*normal and avoiding combinations of antiplatelet therapy ²². Some authors believe that the translaryngeal tracheostomy is superior to other percutaneous techniques in patients with coagulopathy ²³ (Level E).

e. Emergency situations

A patient who has an upper airway obstruction that cannot be relieved by positive pressure mask ventilation or by endotracheal intubation ("cannot intubate, cannot ventilate") must have an immediate surgical airway ²⁴. Although there are several case reports describing successful PT in an emergency situation ²⁵, cricothyrotomy is the method of choice. This is beyond the scope of this protocol.

6) Timing of tracheostomy

The decision when to perform a tracheostomy is controversial ²⁶, although it is known that the number of complications increases after a prolonged duration of endotracheal intubation ²⁷. A frequently cited consensus conference on artificial airways in 1989 recommended endotracheal intubation as the method of choice for an artificial airway needed for up to ten days, whereas tracheostomy is preferred when the need for an artificial airway exceeds 21 days ²⁸ (Level E). A systematic review concluded that performing a tracheostomy at an earlier stage than currently practiced may shorten the duration of artificial ventilation and length of stay in the ICU ²⁹ (Level C). However, this conclusion was drawn from only five studies ³⁰⁻³⁴. Future trials are eagerly awaited. One such study has recently been started in the United Kingdom (www.tracman.org.uk).

Several studies in smaller groups have been performed to assess which patients may profit from an early tracheostomy. In patients with infratentorial lesions ³⁵ or after neurotrauma with a GCS < 7-9 within the first week ^{36;37}, an aggressive policy towards early tracheostomy is justified (Level D). A study prospectively comparing the benefits of early to delayed tracheostomy showed that the benefits of early tracheostomy outweigh the risks of prolonged endotracheal intubation, even in terms of mortality (Level C) ³³. Although this is a very interesting study, there were several unclarified issues ³⁸. For example, the *prediction* as to whether a patient will need more than two weeks of mechanical ventilation is notoriously difficult and often lacks specific objective criteria. Of the many variables during the first 24 hours on the ICU, shock was the only prognostic factor associated with prolonged ventilation (> three weeks) ³⁹. Despite this, 42% of patients with shock on admission were extubated earlier than three weeks. This suggests that including all patients with shock on admission for tracheostomy would be inadvisable.

In conclusion, the decision to convert an endotracheal tube to a tracheostomy canula in the ICU has to be individualized, since firm evidence to support an aggressive approach is lacking. The potential benefits (Table 2) and risks (see 9. Complications) of the procedure compared with prolonging endotracheal intubation need to be considered. Based on the available information, one could consider a tracheostomy as soon as it is apparent that weaning from artificial ventilation is unlikely to happen within two weeks after endotracheal intubation (Level D). The duration of artificial ventilation is shortened when a tracheostomy is placed in an early stage of weaning (Level C). For certain patient categories, in particular neurological patients, the decision to perform a tracheostomy can be made within the first week of admission (Level E).

7) Technique

Like with many other procedures, there is a learning curve with the performance of PT. However, in small ICUs sufficient experience is hard to obtain. It is difficult to define a minimum number of PTs that one needs to perform in order to obtain an acceptable skill level, as this depends on the dexterity of the operator. A reasonable minimum number in the learning phase would be about 15 procedures in order to be proficient to perform the procedure independently ^{40;41} (Level E). It is advisable that a limited number of physicians should be designated to do the procedure ⁴² (Level E). Local circumstances could thus lead to a preference for surgical tracheostomy if training in PT is lacking or caseload is below a certain amount of procedures.

All modern methods for PT rely on the Seldinger technique. Subsequently, dilation up to the degree required for the positioning of the tracheal canula is necessary, either with a single or multiple dilator technique (Table 4)¹. In The Netherlands, almost all ICUs performing PTs use the guide wire dilating forceps or the conic dilation technique ⁴³. In some other countries, there is more experience with the translaryngeal (Fantoni's) technique ^{44;45}. None of the techniques has significant advantages or disadvantages over another and choice depends mainly on individual preference (Level C), although the translaryngeal technique is by far the most complicated.

The preparation of this bedside procedure carried out in the ICU is important. First of all, the patient should be checked for possible contra-indications (Table 3). Patients should be stable, both circulatory and respiratory (for instance, PEEP<10 cm H_2O with PaO₂/FiO₂-ratio>25 kPa) (Level D), although the procedure may be safely done with higher PEEP-levels ⁴⁶. Nasogastric feeding is stopped, the stomach contents are emptied and the hypopharynx sunctioned just before the actual procedure to prevent aspiration of stomach contents into the airway. It is advisable to start ventilating the patients with controlled ventilation with a FiO₂ of 1.0 about 5-10 minutes before the start of the procedure. Care should be taken to compensate for the volume loss due to air leakage during the procedure. Adequate analgesia, sedation and if preferred muscle relaxation should be ensured, according to a local protocol. Local infiltration with lidocaine with epinephrine further reduces the need for analgesia and minimizes bleeding around the incision. Minimal monitoring should be according to the guidelines of the Netherlands Society of Anesthesiologists ⁴⁷, including capnography. The trachea is punctured with a canulated needle attached to a saline or air filled syringe for continuous suction, aiming for the interspace between the first and second or second and third tracheal rings, although one must realize that accurate placement is achieved in less than half of the cases 4^{48} .

The puncture may be guided by fiberoptic view, as this reduces significantly the number of complications compared to PT without bronchoscopy (Level D) ^{15;49}. It helps to confirm the correct position of the puncture, i.e. in the midline of the anterior trachea, and ensures that the posterior wall is not injured. Therefore, arguments that bronchoscopy adds time, cost, and an unnecessary complexity to the procedure and may incur risks to the patient (such as difficulty in maintaining ventilation, CO_2 retention, and elevated intracranial pressures), while true, are weak in comparison to the benefits ¹⁵. An exception can be made for a patient with normal anatomy and an experienced team, but even then a bronchoscope should be readily available in case of unforeseen problems. At least two experienced physicians are required to perform the

procedure safely: one to perform the tracheostomy and one for airway control. An additional assistant may be useful to immobilize the withdrawn tube.

After PT routine chest radiography is unnecessary ⁵⁰ (Level D). The patient with a tracheostomy may be transferred to the general ward, see table 2.7. A removable inner canula should always be used, to facilitate cleaning and to overcome acute canula obstruction. Canula displacement represents a potentially catastrophic complication, in particular in patients who are unable to protect their upper airways (for example EMV<9 or vocal cord paresis) and in particular within the first week of the procedure. Local protocols are important to describe the appropriate aftercare of patients with a tracheostomy canula on the ward, like presence of specific instrumentation (i.e. a tracheal spreader), humidification, physiotherapy, suctioning, stoma care etc. The recent introduction of a consultative intensive care nurse, available 24 hours a day, has also contributed to the care of these patients on the general wards.

8) Airway control during PT

Airway control during PT has several pitfalls, such as the risk of accidental extubation, endotracheal tube cuff rupture, or transfixion of the endotracheal tube. There are several ways to secure the airway, although only the two most relevant methods are discussed here.

a. Tube withdrawal.

One method is to withdraw the endotracheal tube under direct laryngoscopic view prior to puncturing the trachea, so that the cuff is placed in between or just below the vocal cords. It is also possible to withdraw the tube into the pharynx and, following cuff inflation, leaving only the tip into the laryngeal opening, so the tracheal tube cuff acts as a laryngeal inlet obturator.

b. Tube replacement.

It is also possible to replace the endotracheal tube by a laryngeal mask airway (LMA) ⁵¹. However, this method relies on a second technique of airway control, with inherent complications, most importantly aspiration of gastric contents. Intensive care patients often require high inflation pressures, have impaired gastric emptying and have oropharyngeal and perilaryngeal edema secondary to prolonged endotracheal intubation, making emergency re-intubation hazardous. Moreover, bronchoscopy through a LMA is more cumbersome. This method is not advisable ⁵² (Level C).

The best way to deal with any complications during the procedure is to always have at least two physicians available: one to do the procedure and another with experience in airway management for airway control (Level E).

9) Complications of PT

Complications may vary from minor, intermediate to major complications. Minor complications are for example minor perioperative bleeding, mild stomal infection or ugly scarring, while major complications may comprise esophageal perforation, pneumothorax with drainage or tracheal stenosis. Minor complications occur in about 20% of cases, but there is a considerable study-to-study variability in reported complication incidence $(1-58\%)^{53}$. Major complications in PT occur in about 3% (0-14%) and intermediate complications in about 3% (0-26%) of cases 53 . Late complications (after decanulation), although rare, may vary from unesthetic scarring to hoarseness and tracheal stenosis.

Ideally, there should be follow up of patients until the trachea has properly healed for several months after removal of the tracheostomy tube. Unless the events are recorded as critical incidents or as part of an ongoing audit, underreporting of acute complications will occur.

The procedure-related mortality should be defined as mortality associated with the procedure. This mortality rate is less then 0.5% ⁵⁴.

Table 1: Indications for tracheostomy

- 1. Indications for *PT*:
 - a. Any patient who is expected to require mechanical ventilation for at least two weeks with for example:
 - i. Severe (critical illness) polyneuropathy (Level E).
 - ii. Post-multi organ failure, with profound muscle weakness (Level E).
 - iii. Neurological patients with a Glasgow Coma Score < 7-9 and/or an impaired swallow- and cough reflex ³⁵⁻³⁷ (Level D).
 - iv. Severely compromised pulmonary function before admission to the ICU ⁵⁵ (Level D).
 - v. Need for reintubation due to sputum retention. This may also be an indication for a minitracheotomy (see below) (Level E).
 - b. Severe upper airway obstruction (Level E).
- 2. Indication for minitracheotomy. This is limited to:
 - a. Patients where retention of sputum is the only problem 56 (Level D).
- 3. Indications for a primary *surgical* tracheostomy (Level E):
 - a. Expertise for performing a PT not available.
 - b. Patients in need of home ventilation, who require a wide stoma for easy canula changes.
 - c. Patients in whom anatomical landmarks are impossible to localize (although with careful blunt dissection the landmarks may become more clear).
 - d. Oral or nasal intubation impossible or contra-indicated (Level E).

Table 2: Advantages of tracheostomy

- 1. Eating and drinking is possible to some degree (provided the patient is able to swallow) (Level D).
- 2. Speech is possible, whether by deflation of the cuff, or by change of the canula for a fenestrated canula after a <u>minimum of 5 days</u>, in order to let the tracheostomy wound heal sufficiently (Level D).
- 3. Oral hygiene is easier and respiratory secretions are easier removed. The patient is able to cough (Level C).
- 4. Absence of laryngeal and vocal cord injuries. The patient is able to move his or her head more freely and less sedation is needed (Level D).
- 5. Decrease in airway resistance, anatomical dead space and work of breathing, therefore facilitating weaning from mechanical ventilation in patients with marginal respiratory mechanics, although this benefit may be marginal ⁵⁷ (Level E).
- 6. Better security of the airway, because in general a tracheostomy tube can be changed more easily than an endotracheal tube (Level E).
- 7. Depending on local protocols, the patient with a tracheostomy may be transferred to the general ward, provided he/she is able to breathe independently and is able to cough adequately. Tracheal suctioning by nurses should be necessary only once or twice per shift. In the first 48 hours after transfer, close contact with the referring intensive care unit is advisable, for example with the aid of a consulting intensive care nurse.

Table 3: Contra-indications of PT

Absolute:

- 1. Infections at the site of the procedure.
- 2. Uncorrectable coagulation abnormalities.
- 3. Patients in whom anatomical landmarks are impossible to localize.
- 4. Short neck with thyromental distance of less than 3 centimeters, even after optimal exposure.
- 5. Large struma.

Relative:

- 1. Elevated intracranial pressure. Postpone the procedure.
- 2. Emergency situation ("Cannot intubate, cannot ventilate situation").
- 3. Age < 16 years or weight < 40 kg.
- 4. Patients in need of home ventilation, who require a wide stoma for easy canula changes.
- 5. A history of neck surgery and/or irradiation to the neck as the anatomy may be altered.

Technique	Characteristics	References
PDT (Percutaneous dilational	Antegrade, multi-step dilation with up to 7	58;59
tracheostomy)	dilators	
GWDF (Guide wire dilating forceps)	Antegrade, two-step dilation with modified Howard-Kelly forceps	60;61
TLT (Translaryngeal tracheostomy)	Retrograde, single-step dilation with the canula itself	44;45
CDT (Conic dilational tracheostomy)	Antegrade, single-step dilation with a conically shaped, hydrophilically coated dilator	62;63
PercuTwist TM	Antegrade stoma formation with a self- cutting plastic screw	64;65

Table 4. Currently available techniques of percutaneous tracheostomy

Level aanbevelingen

- A. Ondersteund door tenminste twee grote prospectief gerandomiseerde gecontroleerde klinische onderzoeken of een meta-analyse met een kleine kans op een vals positief of een vals negatief resultaat
- B. Ondersteund door één groot prospectief gerandomiseerd gecontroleerd klinisch onderzoek met een kleine kans op een vals positief of een vals negatief resultaat
- C. Ondersteund door één of meerdere kleine prospectief gerandomiseerde gecontroleerde klinische onderzoeken of een meta-analyse met een matige tot grote kans op een vals positief of een vals negatief resultaat
- D. Ondersteund door alleen een niet-gerandomiseerd maar wel gecontroleerd klinisch onderzoek, een cohort studie of een patiëntcontrole onderzoek
- E. Ondersteund door alleen niet-vergelijkend onderzoek, historische controles, case reports of de mening van deskundigen

Reference List

- Staatsen M, Fikkers BG, van der Hoeven JG. Percutaneous tracheostomy. Neth J Crit Care 2005; 9(5):253-258.
- (2) Dongelmans DA, van der Meer NJ, Schultz MJ. [Percutaneous dilatating tracheostomy in intensive-care patients: technique, indications and complications]. Ned Tijdschr Geneeskd 2003; 147(48):2370-2374.
- Al-Ansari MA, Hijazi MH. Clinical review: Percutaneous dilatational tracheostomy. Crit Care 2005; 10(1):202.
- (4) Pierson DJ. Tracheostomy from A to Z: historical context and current challenges. Respir Care 2005; 50(4):473-475.
- (5) Heffner JE, Miller KS, Sahn SA. Tracheostomy in the intensive care unit. Part 1: Indications, technique, management. Chest 1986; 90(2):269-274.
- (6) Goldenberg D, Golz A, Netzer A et al. Tracheotomy: changing indications and a review of 1,130 cases. J Otolaryngol 2002; 31(4):211-215.
- (7) Freeman BD, Isabella K, Lin N et al. A meta-analysis of prospective trials comparing percutaneous and surgical tracheostomy in critically ill patients. Chest 2000; 118(5):1412-1418.
- (8) Delaney A, Bagshaw SM, Nalos M. Percutaneous dilatational tracheostomy versus surgical tracheostomy in critically ill patients: a systematic review and meta-analysis. Crit Care 2006; 10(2):R55.
- (9) Heffner JE. Tracheotomy application and timing. Clin Chest Med 2003; 24(3):389-398.
- (10) Blot F, Melot C. Indications, timing, and techniques of tracheostomy in 152 French ICUs. Chest 2005; 127(4):1347-1352.
- (11) Ben NA, Altman E, Best LA. Extended indications for percutaneous tracheostomy. Ann Thorac Surg 2005; 80(4):1276-1279.
- (12) Blankenship DR, Kulbersh BD, Gourin CG et al. High-risk tracheostomy: exploring the limits of the percutaneous tracheostomy. Laryngoscope 2005; 115(6):987-989.
- (13) Mansharamani NG, Koziel H, Garland R et al. Safety of bedside percutaneous dilatational tracheostomy in obese patients in the ICU. Chest 2000; 117(5):1426-1429.
- (14) Byhahn C, Lischke V, Meininger D et al. Peri-operative complications during percutaneous tracheostomy in obese patients. Anaesthesia 2005; 60(1):12-15.
- (15) Kost KM. Endoscopic percutaneous dilatational tracheotomy: a prospective evaluation of 500 consecutive cases. Laryngoscope 2005; 115(10 Pt 2 Suppl. 107):1-30.
- (16) Curtis JJ, Clark NC, McKenney CA et al. Tracheostomy: a risk factor for mediastinitis after cardiac operation. Ann Thorac Surg 2001; 72(3):731-734.
- (17) Pierce WS, Tyers GF, Waldhausen JA. Effective isolation of a tracheostomy from a median sternotomy wound. J Thorac Cardiovasc Surg 1973; 66(5):841-842.
- (18) Patel NC, Deane J, Scawn N. Reduction in tracheostomy-associated risk of mediastinitis by routine use of percutaneous tracheostomy. Ann Thorac Surg 2002; 73(6):2033.
- (19) Mayberry JC, Wu IC, Goldman RK et al. Cervical spine clearance and neck extension during percutaneous tracheostomy in trauma patients. Crit Care Med 2000; 28(10):3436-3440.
- (20) O'keeffe T, Goldman RK, Mayberry JC et al. Tracheostomy after anterior cervical spine fixation. J Trauma 2004; 57(4):855-860.
- (21) Kluge S, Meyer A, Kuhnelt P et al. Percutaneous tracheostomy is safe in patients with severe thrombocytopenia. Chest 2004; 126(2):547-551.
- (22) <u>www.anesthesiologie.nl/uploads/118/29/Richtlijn_antistolling_DEF.pdf</u>. downloaded January 3th 2006 2006.
- (23) Byhahn C, Lischke V, Westphal K. Translaryngeal tracheostomy in highly unstable patients. Anaesthesia 2000; 55(7):678-682.

- (24) Practice guidelines for management of the difficult airway: an updated report by the American Society of Anesthesiologists Task Force on Management of the Difficult Airway. Anesthesiology 2003; 98(5):1269-1277.
- (25) Ben-Nun A, Altman E, Best LA. Emergency percutaneous tracheostomy in trauma patients: an early experience. Ann Thorac Surg 2004; 77(3):1045-1047.
- (26) Heffner JE. Timing tracheotomy: calendar watching or individualization of care? Chest 1998; 114(2):361-363.
- (27) Marsh HM, Gillespie DJ, Baumgartner AE. Timing of tracheostomy in the critically ill patient. Chest 1989; 96(1):190-193.
- (28) Plummer AL, Gracey DR. Consensus conference on artificial airways in patients receiving mechanical ventilation. Chest 1989; 96(1):178-180.
- (29) Griffiths J, Barber VS, Morgan L et al. Systematic review and meta-analysis of studies of the timing of tracheostomy in adult patients undergoing artificial ventilation. BMJ 2005; 330(7502):1243.
- (30) Bouderka MA, Fakhir B, Bouaggad A et al. Early tracheostomy versus prolonged endotracheal intubation in severe head injury. J Trauma 2004; 57(2):251-254.
- (31) Dunham CM, LaMonica C. Prolonged tracheal intubation in the trauma patient. J Trauma 1984; 24(2):120-124.
- (32) Rodriguez JL, Steinberg SM, Luchetti FA et al. Early tracheostomy for primary airway management in the surgical critical care setting. Surgery 1990; 108(4):655-659.
- (33) Rumbak MJ, Newton M, Truncale T et al. A prospective, randomized, study comparing early percutaneous dilational tracheotomy to prolonged translaryngeal intubation (delayed tracheotomy) in critically ill medical patients. Crit Care Med 2004; 32(8):1689-1694.
- (34) Saffle JR, Morris SE, Edelman L. Early tracheostomy does not improve outcome in burn patients. J Burn Care Rehabil 2002; 23(6):431-438.
- (35) Qureshi AI, Suarez JI, Parekh PD et al. Prediction and timing of tracheostomy in patients with infratentorial lesions requiring mechanical ventilatory support. Crit Care Med 2000; 28(5):1383-1387.
- (36) Gurkin SA, Parikshak M, Kralovich KA et al. Indicators for tracheostomy in patients with traumatic brain injury. Am Surg 2002; 68(4):324-328.
- (37) Major KM, Hui T, Wilson MT et al. Objective indications for early tracheostomy after blunt head trauma. Am J Surg 2003; 186(6):615-619.
- (38) Dongelmans DA, Schultz MJ. Early or late tracheostomy. Crit Care Med 2005; 33(2):466.
- (39) Estenssoro E, Gonzalez F, Laffaire E et al. Shock on admission day is the best predictor of prolonged mechanical ventilation in the ICU. Chest 2005; 127(2):598-603.
- (40) Massick DD, Powell DM, Price PD et al. Quantification of the learning curve for percutaneous dilatational tracheotomy. Laryngoscope 2000; 110(2 Pt 1):222-228.
- (41) Liao L, Myers J, Johnston J et al. Percutaneous tracheostomy: one center's experience with a new modality. Am J Surg 2005; 190(6):923-926.
- (42) Petros S. Percutaneous tracheostomy. Crit Care (Lond) 1999; 3(2):R5-R10.
- (43) Fikkers BG, Fransen GA, van der Hoeven JG et al. Tracheostomy for long-term ventilated patients: a postal survey of ICU practice in The Netherlands. Intensive Care Med 2003; 29(8):1390-1393.
- (44) Fantoni A, Ripamonti D. A non-derivative, non-surgical tracheostomy: the translaryngeal method. Intensive Care Med 1997; 23(4):386-392.
- (45) Sharpe MD, Parnes LS, Drover JW et al. Translaryngeal tracheostomy: experience of 340 cases. Laryngoscope 2003; 113(3):530-536.
- (46) Beiderlinden M, Groeben H, Peters J. Safety of percutaneous dilational tracheostomy in patients ventilated with high positive end-expiratory pressure (PEEP). Intensive Care Med 2003; 29:944-948.

- (47) <u>www.anesthesiologie.nl/verenigingsstandpunten?wid=169&func=viewSubmission&sid=245&pageld=9</u>. downloaded January 3th 2006 2006.
- (48) Hotchkiss KS, McCaffrey JC. Laryngotracheal injury after percutaneous dilational tracheostomy in cadaver specimens. Laryngoscope 2003; 113(1):16-20.
- (49) Blankenship DR, Gourin CG, Davis WB et al. Percutaneous tracheostomy: don't beat them, join them. Laryngoscope 2004; 114(9):1517-1521.
- (50) Hoehne F, Ozaeta M, Chung R. Routine chest X-ray after percutaneous tracheostomy is unnecessary. Am Surg 2005; 71(1):51-53.
- (51) Dexter TJ. The laryngeal mask airway: a method to improve visualisation of the trachea and larynx during fibreoptic assisted percutaneous tracheostomy. Anaesth Int Care 1994; 22(1):35-39.
- (52) Ambesh SP, Sinha PK, Tripathi M et al. Laryngeal mask airway vs endotracheal tube to facilitate bedside percutaneous tracheostomy in critically ill patients: a prospective comparative study. J Postgrad Med 2002; 48(1):11-15.
- (53) Fikkers BG. Percutaneous Tracheostomy on the Intensive Care unit. Thesis 2004.
- (54) McCormick B, Manara AR. Mortality from percutaneous dilatational tracheostomy. A report of three cases. Anaesthesia 2005; 60(5):490-495.
- (55) Boynton JH, Hawkins K, Eastridge BJ et al. Tracheostomy timing and the duration of weaning in patients with acute respiratory failure. Crit Care 2004; 8(4):R261-R267.
- (56) Bonde P, Papachristos I, McCraith A et al. Sputum retention after lung operation: prospective, randomized trial shows superiority of prophylactic minitracheostomy in high-risk patients. Ann Thorac Surg 2002; 74(1):196-202.
- (57) Diehl JL, El Atrous S, Touchard D et al. Changes in the work of breathing induced by tracheotomy in ventilator-dependent patients. Am J Resp Crit Care Med 1999; 159(2):383-388.
- (58) Ciaglia P, Firsching R, Syniec C. Elective percutaneous dilatational tracheostomy. A new simple bedside procedure; preliminary report. Chest 1985; 87(6):715-719.
- (59) Wang MB, Berke GS, Ward PH et al. Early experience with percutaneous tracheotomy. Laryngoscope 1992; 102(2):157-162.
- (60) Griggs WM, Worthley LI, Gilligan JE et al. A simple percutaneous tracheostomy technique. Surg Gynecol Obstet 1990; 170(6):543-545.
- (61) Fikkers BG, van Heerbeek N, Krabbe PF et al. Percutaneous tracheostomy with the guide wire dilating forceps technique: presentation of 171 consecutive patients. Head Neck 2002; 24(7):625-631.
- (62) Byhahn C, Lischke V, Halbig S et al. [Ciaglia Blue Rhino: a modified technique for percutaneous dilatation tracheostomy. Technique and early clinical results]. Anaesthesist 2000; 49(3):202-206.
- (63) Fikkers BG, Briede IS, Verwiel JM et al. Percutaneous tracheostomy with the Blue Rhino technique: presentation of 100 consecutive patients. Anaesthesia 2002; 57(11):1094-1097.
- (64) Frova G, Quintel M. A new simple method for percutaneous tracheostomy: controlled rotating dilation. A preliminary report. Intensive Care Med 2002; 28(3):299-303.
- (65) Sengupta N, Ang KL, Prakash D et al. Twenty months' routine use of a new percutaneous tracheostomy set using controlled rotating dilation. Anesth Analg 2004; 99(1):188-192.